

## TESTING THE LIMITS OF INTEGRATION TIME (AND OBSERVER PATIENCE): DETECTION OF INTERSTELLAR KCN

C. SAVAGE, L.M. ZIURYS, *Department of Chemistry, Department of Astronomy, and Steward Observatory, University of Arizona, Tucson, AZ 85721*; M. GUÉLIN, *Institut de RadioAstronomie Millimétrique, 300 rue de la Piscine, Domaine Universitaire, 38406 Saint Martin d'Hères, France*.

KCN has been detected in the circumstellar envelope of IRC+10216, a late-type carbon star, with the Kitt Peak 12-meter radio telescope. KCN is a T-shaped, closed shell molecule with a typical asymmetric top spectrum. Five  $K_a = 0$  components were detected in the  $\lambda = 2\text{-}3.5$  mm wavelength range. All five lines were fit to a linear rotational diagram, indicating that the KCN emission results from a uniform rotational temperature of 27 K. The derived column density of KCN is  $\sim 1 \times 10^{12}$  cm $^{-2}$ , corresponding to a fractional abundance of  $\sim 3 \times 10^{-10}$  relative to H $_2$ . All of the lines observed had intensities of 1-2 mK, making them the weakest yet detected by mm-wave astronomy. In order to achieve  $3\sigma$  detections, integration times for all lines were at least 50 hours, with some frequencies requiring over 100 hours to attain sufficient signal-to-noise. These observations demonstrate the stability of the 12-meter over long periods of time, and confirm the theoretical noise behavior as given by the radiometer equation:

$$T_{rms} = \frac{2T_{sys}}{\eta_{spec} \sqrt{\Delta\nu t_{scan}}}$$